References/ Bibliography:

- Aliero, B.L (2003). Larvicidal effects of aqueous extracts of Azardirachta indica (neem) on the larvae of Anopheles mosquito, African Journal of biotechnology, 2 (9), pp 325-327.
- Anitha, V., Rogers, D.J., Wightman, J. and Ward, A. (2006). Distribution and abundance of white grubs (Coleoptera: Scarabaeidae) on groundnut in southern India. Crop Protection, 25, 732–740.
- Asmat M, Kangoo Zakir, H Khan and Shahnawaz Ahmad (2011). External Morphology of Holotrichia consanguinea Blanchard (Coleoptera, Scarabaeidae, Melolonthinae). Research Journal of Agricultural Sciences, 2(1): 146-147.
- Beatrice Kaufmann and Philippe Christen (2002). Recent Extraction Techniques for Natural Products: Microwave-assisted Extraction and Pressurised Solvent Extraction. Phytochem. Anal. 13, 105–113.
- Benigno G., Adriana C., Christiane J., Lorena R.M., Francisco J. V. (2000). Ethnoecology of White Grubs (Coleoptera: Melolonthidae) among the Tzeltal maya of Chiapas. Journal of Elhnobiology 20(1): 43-59.
- Bhanot, R. K. & G. P. Bhawane (1991). Effect of plant extractives on intestinal phosphatases of the adult, H. Serrata (Coleoptera: Scarabaeidae) J. Curr. Biosci., 8: 39-44.
- C. Thamarai chelvi, W. Richard Thilagaraj, R. Kandasamy (2010). Laboratory culture and virulence of Beauveria brongniarti (Metschinikoff) isolates on sugarcane white grub, Holotrichia serrata (Coleoptera : Scarabidae) Journal of Biopesticides 3(1 Special Issue) 177 – 179.

- Chandel, R.S., Gupta, P.R. and Thakurm, J.R., (1997). Host preference and seasonal abundance of defoliating beetles infesting fruit trees in mid hills of Himachal Pradesh. J. Soil Biol. and Ecol. 17(1): 140-146.
- Chandel, R.S., Chandla, V.K. and Ashwani Sharma (2003). Population dynamics of potato white grubs in Shimla hills. J. Potato Assoc. 30(1-2): 151-152.
- Chirame, B.B., Khadatare, R.M. and Bhoi, P.G., (2003). Control of white grub (Holotrichia serrata F.) in sugarcane with Beauveria brongniartii. J. Maharashtra Agric. Uni. 28(2): 208-209.
- D. S. Nikam, S. D. Disale, R. G. Kudale & G. P.Bhawane (1997). Comparative study on the haemocyte of scarabaeidae, Indian J. Comp.Anim. Physiol (15): 27-30.
- David J. Shetlar (2001). Control of Japnese beetle adults and grubs in home lawns, Extension Fact Sheet, HYG- 03.
- G. P. Bhawane & R. K. Bhanot (1992). Impact of different plant extractive on the digestive carbohydrases of the adult Leucopholis lepidophora Bl. J. Adv. Zool. 13 (1& 2): 19-24.
- G. P. Bhawane, et al… (2011). Life cycle of *Holotrichia karschi* brenske (coleoptera: scarabaeidae: melolonthinae), The Bioscan 6(3), 771-774.
- Garg, V.K., Kulkarni, N. and Meshram, P.B. (2005). Influence of weather parameters on the incidence of white grub, Holotrichia serrata Fabr. on teak seedlings in Chhindwara Forest Divisions. Indian Journal of Tropical Biodiversity, 13, 48–51.
- Gupta, B.M. and Yadava, C.P.S., (1993). Distribution pattern of white grub Holotrichia consanguinea Blanch in semi arid Rajasthan. Indian J. Ent. 55(1): 314-316.

- J. A. Wightman and G.V. Ranga Rao (1993). A Groundnut Insect Identification Handbook for India. ICRISAT Information Bulletin no. 39 International Crops Research Institute for the Semi -Ar id Tropics Patancheru, Andhra Pradesh 502 324, India
- Jasna I., Irena Z., Slobodan P., Dejan S. (2009). The analysis of different processes of extraction: yield of extracts obtained from Aloe vera (Aloe barbadensis miller) and sweet bay (Laurus nobilis l.) and the energy analysis of applied processes. Chemical Industry & Chemical Engineering Quarterly 15 (4) 271−278.
- Meshram, P.B., Joshi, K.C. and Sarkar, A.K. (1993). Efficacy of some insecticides against the white grubs, Holotrichia insularis Brenske in teak nursery. Annals of Forestry, 1, 196–198.
- Mishra, P.N. and Singh, M.P., (1993). Determination of predominant species of white grub in Kumaon region of UP hills and their control. Ann. Pl. Prot. Sci. 5(2): 154-156.
- Musthak Ali, T.M. (2001). Biosystematics of phytophagous Scarabaeidae – an Indian overview. Indian Phytophagous Scarabaeidae and Their Management (eds. G. Sharma, Y.S. Mathur & R.B.L. Gupta), pp. 5–47. Agrobios, India.
- Nandagopal V and Prasad TV. (2004). World list of insect and non insect pests of groundnut and their natural enemies. Junagadh, Gujarat, India: National Research Centre for Groundnut. pp. 10–14.
- Patil SB, Wagh SR, Sakate SD and Bhawane GP (2006). Effect of insecticides on the digestive enzymes of adults Holotrichia serrata. Indian J. Comp. Anim. Physiol. Vol. 24(1) p 47-51.

- Patil, S.M. and Adsule, V.M., (1991). Population density of different stages of Leucopholis lepidophora Blanch infesting sugarcane (Coleoptera : melolonthinae). Indian Sug. 41(1): 169-171.
- Procedures for soil analysis (2002). International Soil Reference and Information Centre, ISBN: 90-6672-044-1
- Ranga Rao GV, et al… (2006). Occurance of White Grubs in Groundnut Crop in Uplands of South Vietnam: A new report.
- Recommended Chemical Soil Test Procedures for the North Central Region (2011). North Central Regional Research Publication No. 221 (Revised).
- S.B. Patil, S.R. Wagh, S.D. Disale and G. P. Bhawane (2005). Effect of different insecticides on fecundity and egg hatchability in Holotrichia serrata Fab. (Coleoptera: Scarabaediae). Indian J. Comp. Anim. Physiol. 23(1): 41-44.
- S. D. Disale, G. P.Bhawane, N. K. More & B. K. Pawar (1993). Alternations in Blood volume and total haemocyte count and utilization of dissolved oxygen by the fossorial grubs of Leucopholis lepidophora Bl. (Coleoptera: Scarabaeidae). Comp. Physiol. Ecol., 18(4): 131-135.
- S. Karuppusamy and K.M. Rajasekaran (2009). High Throughput Antibacterial Screening of Plant Extracts by Resazurin Redox with Special Reference to Medicinal Plants of Western Ghats. Global Journal of Pharmacology, 3 (2): 63-68.
- Shambhukar DA, et al… (2006) Assesment of Integrated pest management modules in Groundnut on Farmers fields. IAN 26.
- Sri Wardani, Sugiyarto (2009). Characterization of white grubs (Melolonthidae: Coleoptera) at salak pondoh agroecosystem in Mount Merapi based on isozymic banding patterns Nusantara Bioscience, Vol. 1, No. 1, Pp. 38-42.

- Singh, M.P. and Mishra, P.N., (2003). White grubs in hilly and mountain agro ecosystem of Uttaranchal. Indian Farm, 1(1): 20-23.
- Singh R.P., (1993). In Botanical Pesticides in Developing Countries: Current Status and Future Trends (Eds Dhaliwal, G.S. and Singh, B.Coomon Wealth Publishers, New Delhi, pp.236-268.
- Wightman, J.A. and Ranga Rao, G.V. (1993). A Groundnut Insect Identification Handbook for India. Information Bulletin no. 39. (In En. Summaries in En, Fr.) Patancheru, A.P. 502 324, India: International Crops Research Institute for the Semi-Arid Tropics. 64 pp.
- Wightman, J.A., Ranga Rao, G.V., (1994). Groundnut pests. In: Smartt, J. (Ed.), the Groundnut Crop: a Scientific Basis for Improvement. Chapman & Hall, London, pp. 395–469.
- Yadava, C.P.S. and Sharma, G.K. (1995). Indian white grubs and their management. All India Coordinated Research Project on white grubs, Technical Bulletin No. 2, ICAR, New Delhi, India, 26 pp.